


CIRCULATION COPY  
SUBJECT TO REVIEW  
BY THE AUTHOR

## Demonstration of a Soft X-ray Amplifier

D. Matthews, P. Hagelstein, M. Rosen,  
M. Eckart, N. Ceglio, A. Hazi, H. Medeck  
B. McGowan, J. Trebes, B. Whitten,  
E. Campbell, C. Hatcher, A. Hawryluk  
R. Kauffman, L. Pleasance, G. Rambach,  
J. Scofield, G. Stone and T. Weaver

Physics Review Letters



This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

This work performed under the auspices  
of the U.S. Department of Energy by  
the Lawrence Livermore National Laboratory  
under contract #W-7405-ENG-48.

#### **DISCLAIMER**

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial products, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

## Demonstration of a Soft X-ray Amplifier

D. L. Matthews, P. L. Hagelstein, M. D. Rosen, M. J. Eckart,  
N. M. Ceglio, A. U. Hazi, H. Medeck, B. J. MacGowan,\*  
J. E. Trebes, B. L. Whitten, E. M. Campbell,  
C. W. Hatcher, A. M. Hawryluk, R. L. Kauffman, L. D. Pleasance,  
G. Rambach, J. H. Scofield, G. Stone, and T. A. Weaver

University of California  
Lawrence Livermore National Laboratory  
Livermore, California 94550

### Abstract

We report observations of amplified spontaneous emission at soft x-ray wavelengths. An optical laser ionized thin foils of selenium to produce a population inversion of the  $2p^5 3p$  and  $2p^5 3s$  levels of the neon-like ion. Using three time-resolved, spectroscopic measurements we demonstrated gain-length products up to 6.5 and gain coefficients of  $5.5 \pm 1.0 \text{ cm}^{-1}$  for the  $J=2$  to 1 lines at 206.3 and 209.6 Å. We also observed considerable amplification for the same transitions in yttrium at 155.0 and 157.1 Å.

\*Permanent address: Blackett Laboratory, Imperial College, London, SW7 2AZ, England.

PACS Numbers: 42.60.By, 42.55.-f, 32.30.Rj, 32.70.-n

For over a decade there has been an intense search for a way to produce amplification of XUV or soft x-ray radiation. Several recent papers<sup>1,2</sup> review the progress of the theoretical and experimental efforts. In general, the literature lacks a conclusive or unrefuted demonstration of such amplification.

One of the proposed methods of producing an XUV amplifier uses a high-power optical laser to directly heat a plasma. Using this technique, Zherikhin, Koshelev, and Letokhov<sup>3</sup> first described a mechanism for obtaining an inversion between  $2p^5 3p$  and  $2p^5 3s$  levels in neon-like ions. Vinogradov and co-workers<sup>4</sup> have subsequently written a series of papers refining the theoretical description. To produce the inversion, the  $n=3$  excited levels are populated by electron impact excitation from the ground ( $2p^6$ ) state of the neon-like ion, which is itself produced in the plasma heated by the optical laser. The population inversion between  $2p^5 3p$  and  $2p^5 3s$  levels develops because of the large difference between the radiative decay rates. Recent calculations<sup>5-9</sup> have extended this neon-like excitation scheme to ions in which the lasing wavelengths approach the soft x-ray regime.

The purpose of this letter is to report the first results from new experiments on the neon-like collisional excitation scheme. We describe the first conclusive demonstration of a macroscopic-sized gain medium which exhibits substantial amplification of at least four  $2p^5 3p-2p^5 3s$  transitions in selenium, with the largest being observed for the  $J=2$  to  $1$  lines at 206.3 and 209.6 Å. The transition with the largest predicted gain,<sup>4-9</sup> the  $J=0$  to  $1$  transition at approximately 183 Å, has not been

uniquely identified. We base our conclusions on the results of over 100 separate laser-irradiated target experiments. We used a variety of spectroscopic instruments to measure the time-resolved absolute brightness of the amplified spontaneous emission (ASE), and to demonstrate the non-linear increase of the emission with increased length of the gain medium. The anisotropy of the ASE was also verified by measuring the relative intensity in the axial direction and in another direction far off that axis.

These x-ray laser experiments were conducted at the Novette laser-target irradiation facility. A simplified schematic representation of the experimental arrangement is shown in Fig. 1. The exploding foil target design<sup>10</sup> is based on experience with long density-scale-length plasmas that are produced in inertial confinement fusion research. Typically, the target was composed of a 750 Å layer of Se, vapor deposited on one side of a 1500 Å-thick Formvar substrate. The foil was nominally 1.1 cm-long and was illuminated by green light ( $\lambda = 0.532 \mu\text{m}$ ) along a line focus with dimensions of 0.02x1.12 cm. The nominal pulse length was 450 psec, and the typical incident intensity was  $5 \times 10^{13} \text{ W/cm}^2$ . The targets were irradiated in two different geometries: "single-sided" in which a given segment of the foil was hit by only one laser beam, and "double-sided" in which opposing laser beams irradiated a common target area. Using the single-sided geometry we were able to illuminate targets up to 2.2 cm long by displacing the two beams axially.

X-ray emission in the two axial directions (see Fig. 1) was measured using a grazing incidence spectrograph (GIS) and a transmission grating spectrograph (TGSS). The GIS has high spectral resolution ( $\lambda/d\lambda = 1800$ ) and a line radiation detection threshold of 6 erg/sr. It utilizes a microchannel plate contoured to the Rowland circle for detection of the x-ray spectra. Gating the microchannel plate detector by an Auston switch allowed viewing of the spectrum for a limited period of time (250-750 psec), thereby discriminating against emission occurring late in the experiment. The TGSS has a spectral resolution of  $\lambda/d\lambda = 200$ , detection threshold of 3.9 mW/Å (radiated into a solid angle of  $\sim 1.1 \times 10^{-4}$  sr), and temporal resolution of 20-30 psec. The instrument incorporates an ellipsoidal collection optic, a 2000 Å period transmission grating and a soft x-ray streak camera. In addition to the emission along the target's longitudinal axis, we recorded the spectrum at  $77^\circ$  off-axis using another GIS which viewed the entire target.

In the course of the experiments we measured strong anisotropy of certain lines emitted from the exploding foils. Figures 2a,b illustrate spectral data monitored by the two GIS in the same time segment. Note the presence of Na-like resonance transitions (not expected to be amplified) both on and off axis, whereas the strong lines at 206.3 and 209.6 Å are only seen on axis. Based on ab initio calculations and isoelectronic extrapolation<sup>11</sup> from lower Z, these lines are positively identified as the  $(2p_{3/2}3p_{3/2})_{J=2} - (2p_{3/2}3s)_{J=1}$  and  $(2p_{1/2}3p_{3/2})_{J=2} - (2p_{1/2}3s)_{J=1}$  transitions of Ne-like Se,

respectively. With the exception of the  $3d_{3/2} - 3p_{3/2}$  transition at 178.6 Å, the Na-like lines are optically thick and appear brighter off-axis because the GIS views a larger surface area of the plasma in that direction.

In Figure 2c we plot a theoretical spontaneous emission spectrum of Ne-like transitions calculated under the assumption of an optically thin plasma in steady state at  $T_e = 1.0$  keV and  $N_e = 1 \times 10^{21} \text{ cm}^{-3}$ . In the experimental spectra, the 206.3 and 209.6 Å lines are much stronger than all of the other non-lasing neon-like transitions which have larger gf-values, e.g. the  $2p^5 3d - 2p^5 3p$  lines at 191.5, 188.4, and 169.2 Å (see Ref. 11). The dominance of the two J=2 to 1 lines in the spectrum was observed in approximately 100 laser shots.

Figure 2d illustrates the maximum amplification we obtained in these experiments. In this case the target was shot single-sided at an incident intensity of  $1 \times 10^{14} \text{ W/cm}^2$ , but with the beams displaced to give a total amplification length of 2.2 cm. It is important to note that we do not observe a strong line near 183 Å, the predicted position of the transition with the largest calculated<sup>4-9</sup> gain  $[(2p_{1/2} 3p_{1/2})_{J=0} - (2p_{1/2} 3s)_{J=1}]$ . If there were little or no amplification of this transition, it could not be seen because of its small gf value.<sup>5,8,11</sup>

Figure 3 illustrates the temporal behavior of the emission as recorded by the TGSS. Figure 3(a) shows film data obtained with a 1.0 cm target displaying moderate amplification. In Fig. 3(b) the temporal profile of the amplified line at 209.6 Å is compared with that of the spontaneous emission of the Na-like ion at 201.1 Å. Note that the time

duration of the amplified emission is significantly shorter than that of the nearby spontaneous lines as well as the pulsewidth of the pump laser. With increased amplification, the pulsewidths of the lasing lines tend to decrease. Figure 3(c) shows film data obtained with a 2.2 cm target for which the amplification is large. The insert shows a spectral profile taken at the time of peak ASE. The width of the line, which is source-size broadened, is  $\sim 1.0 \text{ \AA}$  indicating a source region of  $\sim 200 \text{ }\mu\text{m}$  diameter. Figure 3(c) gives a strong indication of non-equilibrium conditions in the plasma. If we assume that the  $206.3 \text{ \AA}$  line is Doppler-broadened, with a width of  $0.04 \text{ \AA}$  ( $T_{\text{ion}} = 400 \text{ eV}$ ), then we obtain  $\sim 40 \text{ keV}$  for the equivalent brightness temperature. In contrast, the brightness temperature inferred from the Na-like emission at  $201.1 \text{ \AA}$  is  $\sim 0.1 \text{ keV}$ .

The most quantitative evidence for amplification is shown in Fig. 4. These results come from the axial GIS and are integrated over the linewidths of the  $206.3$  and  $209.6 \text{ \AA}$  transitions. The data exhibit a non-linear dependence on the length of the amplifier. The output intensity (integrated over frequency) emitted by a one-dimensional homogeneous gain medium of length  $L$  under steady state conditions scales as  $\exp(\alpha L)L^{-1/2}$ , where  $\alpha$  is the line center gain coefficient. This scaling is valid when the output divergence is greater than the detector solid angle and  $\alpha L \geq 2$ . Fitting the data from double-sided shots to this expression yields a gain coefficient of  $\alpha = 5.5 \pm 1.0 \text{ cm}^{-1}$  for both the  $206.3$  and  $209.6 \text{ \AA}$  lines. Although the data obtained with single-sided irradiation demonstrates significant amplification, up to



700 times spontaneous levels, the complexity of both the target irradiation geometry and the radiation transport preclude simple modeling of the length dependence. Note also that the behavior of the data as a function of length indicates the amplifier is below saturation.

In addition to the measurements described above, we have done preliminary experiments to produce amplification at shorter wavelengths. In particular, we irradiated yttrium foil targets and observed amplification of the  $J=2$  to 1 neon-like transitions at 155.0 and 157.1 Å. Once again, the neon-like  $J=0$  to 1 transition was not observed. Just as for selenium, the yttrium emission exhibited strong anisotropy, short time duration, and a brightness temperature of  $\sim 5$  keV (155.0 Å line) .

In conclusion, by using an optical laser to produce a population inversion in a simple exploding-foil, we have demonstrated substantial amplification of spontaneous emission at soft x-ray wavelengths.

The authors gratefully acknowledge specific contributions from T. Barbee, B. Boyd, D. Christie, D. Dietrich, M. Gerassimenko, G. Heaton, S. Hildum, G. Howe, K. Manes, D. Nilson, G. Power, L. Seppala, R. Stewart, S. Stribling, J. Tassano, A. Toor, J. Trenholme, G. Vayer, and J. Wiedwald. This work performed under the auspices of the U. S. Department of Energy by the Lawrence Livermore National Laboratory under contract #W-7405-ENG-48.

References


1. R. C. Elton, Adv. in X-ray Analysis 21, Plenum, New York (1978) p. 1.
2. F. V. Bunkin, V. I. Derzhiev, and S. I. Yakovlenko, Sov. J. Quant. Elect. 11, 981(1981).
3. A. Zherikhin, K. Koshelev, and V. Letokhov, Sov. J. Quant. Mech. 6, 82(1976).
4. A. V. Vinogradov and V. Shlyaptsev, Sov. J. Quant. Elect. 13, 1511(1983), and references therein.
5. U. Feldman, A. Bhatia, and S. Suckewer, J. Appl. Phys. 54, 2188(1983);  
U. Feldman, J. F. Seely, and A. K. Bhatia, J. Appl. Phys. 56 2475 (1984).
6. J. P. Apruzese and J. Davis, Phys. Rev. A28, 3686(1983).
7. G. Dahlbacka, R. Dukart, R. Fortner, D. Dietrich, and R. Stewart, 1982 IEEE International Conference on Plasma Science, Carlton University, Ottawa, Canada, May 17-19, 1982. p. 39.
8. R. Stewart, Ph.D. thesis, Lawrence Livermore National Lab Report, UCRL\_\_\_\_ (1984).


9. P. Hagelstein, Plasma Phys 25, 1345(1983).
10. M. D. Rosen, et al, preceding paper.
11. M. Eckart, J. Scofield, A. Hazi private communication.

#### Figure Captions


1. Placement of primary diagnostics used in this experiment.
2. (a) Grazing incidence spectrograph (GIS) data from off-axis line-of-sight. Data taken from beginning of radiative output until 500 psec after peak of pump laser. (b) On-axis GIS data from same time interval as 2a. (c) Calculated spontaneous spectrum of neon-like selenium. (d) On-axis GIS data from maximum gain target.
3. (a) Film data from the TGSS for a moderate gain target. (b) Temporal profiles of the neon-like 209.6 Å lasing line (A) and the Na-like 201.1 Å line (B). (c) Film data from TGSS for a maximum gain target. Inset shows power vs. wavelength.
4. Integrated line intensity of  $J=2$  to 1 transitions vs. amplifier length for (a) double-sided and (b) single-sided laser irradiation conditions.







(a)



(b)

